arXiv:1608.01550v1 [math.GR] 4 Aug 2016

Asymmetric dynamics of outer automorphisms

Mark C. Bell
University of Illinois
mcbell@illinois.edu

August 5, 2016

Abstract

We consider the action of an irreducible outer automorphism ϕ on the closure of Culler–Vogtmann Outer space. This action has north-south dynamics and so, under iteration, points converge exponentially to $[T_+^\phi]$.

For each $N \geq 3$, we give a family of outer automorphisms $\phi_k \in \mathrm{Out}(\mathbb{F}_N)$ such that as, k goes to infinity, the rate of convergence of ϕ_k goes to infinity while the rate of convergence of ϕ_k^{-1} goes to one. Even if we only require the rate of convergence of ϕ_k to remain bounded away from one, no such family can be constructed when $N < 3$.

This family also provides an explicit example of a property described by Handel and Mosher: that there is no uniform upper bound on the distance between the axes of an automorphism and its inverse.

keywords. Outer automorphism, rate of convergence, asymmetric.

Mathematics Subject Classification 2010: 37E25, 20E05, 37D20

An irreducible outer automorphism $\phi \in \mathrm{Out}(\mathbb{F}_N)$ acts on $\overline{CV_N}$, the closure of Culler–Vogtmann Outer space [11, Section 1.4], with north-south dynamics [8, Theorem 1.1]. Therefore its action has a pair of fixed points $[T_+^\phi], [T_-^\phi] \in \partial CV_N$ and under iteration points in $\overline{CV_N}$ (other than $[T_-^\phi]$) converge to $[T_+^\phi]$.

There is a natural embedding of $\overline{CV_N}$ into $\mathbb{RP}^{\mathcal{C}}$, where $\mathcal{C}$ is the set of conjugacy classes of the free group $\mathbb{F}_N$. In this embedding, the $w \in \mathcal{C}$ coordinate of a marked graph $(G, g) \in CV_N$ is given by the length of the shortest loop in G which is freely homotopic to $g(w)$ [11, Page 5]. In this coordinate system the convergence to $[T_+^\phi]$ is exponential. To measure the rate of this convergence we recall two definitions from [1].

Definition 1 ([1, Definition 1.2])**.** Suppose that $f \in \mathbb{Z}[x]$ is a polynomial with roots $\lambda_1, \ldots, \lambda_m$, ordered such that $|\lambda_1| \geq |\lambda_2| \geq \cdots \geq |\lambda_m|$. The *spectral ratio* of f is $\omega(f) := |\lambda_1/\lambda_2|$.

Definition 2. Suppose that $\phi \in \mathrm{Out}(\mathbb{F}_N)$ is an irreducible outer automorphism. Let $\mu_{\lambda(\phi)} \in \mathbb{Z}[x]$ denote the minimal polynomial of its stretch factor $\lambda(\phi)$. The *spectral ratio* of ϕ is $\omega(\phi) := \omega(\mu_{\lambda(\phi)})$.

As described in [1], the rate of convergence of $[T \cdot \phi^n]$ to $[T^{\phi}_{+}]$ is determined by $\omega(\phi)$. Thus we state the main result of this paper, a complete characterisation of when it is possible to build outer automorphisms which converge rapidly in one direct but slowly in the other, in terms of the spectral ratio:

Theorem 3. *There is a family of fully irreducible outer automorphisms $\phi_k \in \mathrm{Out}(\mathbb{F}_N)$ such that $\omega(\phi_k) \to \infty$ but $\omega(\phi_k^{-1}) \to 1$ if and only if $N \geq 3$.*

This gives another difference between irreducible outer automorphisms and pseudo-Anosov mapping classes of surfaces. A pseudo-Anosov mapping class $h \in \mathrm{Mod}^+(S)$ has its spectral ratio $\omega(h)$ defined in terms of its dilatation [1, Definition 1.3]. However, h and h^{-1} have the same dilatation [3, Proposition 11.3] and so $\omega(h) = \omega(h^{-1})$ automatically.

It is straightforward to prove the forward direction of Theorem 3 by considering its contrapositive:

- When $N = 1$ there is nothing to check as all automorphisms of $\mathbb{F}_1$ are finite order.
- When $N = 2$, any irreducible outer automorphism $\phi \in \mathrm{Out}(\mathbb{F}_2)$ is geometric. Hence there is a pseudo-Anosov mapping class on $S_{1,1}$, the once-punctured torus, which induces ϕ on $\pi_1(S_{1,1})$.

  However, the spectral ratio of any pseudo-Anosov mapping class of the once-punctured torus is at least $\varphi^4 = 6.854101\cdots$, where φ is the golden ratio [1, Section 3.3]. Therefore $\omega(\phi) = \omega(\phi^{-1}) \geq \varphi^4$ and so these are both bounded away from one.

Thus we devote the remainder of this paper to constructing an explicit family of outer automorphisms of $\mathbb{F}_N = \langle a_1, a_2, \ldots, a_N \rangle$, for fixed $N \geq 3$.

To do this, we start by considering the polynomials

$$f(x, y) := x^N - yx^{N-1} - 1 \quad \text{and} \quad g(x, y) := x^N - yx^{N-2} - 1$$

in $\mathbb{Q}[x, y]$. Since these are linear polynomials in y, they are irreducible over $\mathbb{Q}$. Therefore, by Hilbert's irreducibility theorem [7, Chapter 9], there are infinitely many integers $k \geq 3$ such that $p_k(x) := f(x, k)$ and $q_k(x) := g(x, k)$ are both irreducible.

Lemma 4. *Suppose that $k \geq 3$. The polynomial p_k has $N - 1$ roots inside of the unit circle and one root in $[k, k+1]$. Similarly, the polynomial q_k has $N-2$ roots inside of the unit circle, one root in $[-\sqrt{k+1}, -\sqrt{k-1}]$ and one root in $[\sqrt{k-1}, \sqrt{k+1}]$. For example, see Figure 1.*

Proof. Let $h(z) := -kz^{N-1}$. Then when $|z| = 1$ we have that

$$|(p_k - h)(z)| = |z^N - 1| \leq 2 < |p_k(z)| + |h(z)|.$$

Therefore, by Rouché's theorem, p_k must have the same number of roots inside of the unit circle as h does. Hence p_k has $N - 1$ roots inside of the unit circle.

Furthermore, since $\deg(p_k) = N$, there is only one more root to locate. The intermediate value theorem now shows it must lie in $[k, k + 1]$.

Applying the same argument with $h(z) := -kz^{N-2}$ to q_k shows it must have $N - 2$ roots inside of the unit circle. Again, we verify that the other two roots of q_k lie in $[-\sqrt{k+1}, -\sqrt{k-1}]$ and $[\sqrt{k-1}, \sqrt{k+1}]$ by using the intermediate value theorem. □



Figure 1: The polynomials p_{10} and q_{100} when $N = 3$.

In fact knowing the positions of the roots allows us to show many of these polynomials are irreducible directly.

Lemma 5. *Whenever $k \geq 3$, the polynomial p_k is irreducible over $\mathbb{Z}$.*

Proof. First assume that there is a z such that $|z| = 1$ and $p_k(z) = 0$ and so $1 = |z^N - kz^{N-1}| = |z - k|$. However this would mean that $|k| \leq |k - z| + |z| = 2$ which contradicts the fact that $k \geq 3$. Hence p_k cannot have a root on the unit circle.

Now assume that p_k is reducible. Then one of its factors, $q \in \mathbb{Z}[x]$, must have all of its roots inside of the unit circle. Therefore the constant term of q, which is the product of its roots, must have modulus less than one. However this means that the constant term of q must be zero and so zero is a root of p_k, which is false. □

Similarly, by taking into account the symmetry of the roots, the same argument also shows that q_k is irreducible whenever $k \geq 3$ and N is even. Furthermore, for many of the low, odd values of N we can find a prime p and $k' \in \mathbb{Z}/p\mathbb{Z}$ such that the image of $q_{k'}$ in $(\mathbb{Z}/p\mathbb{Z})[x]$ is irreducible. It then follows that q_k is irreducible whenever $k \equiv k' \pmod{p}$. Some of these values are shown in Table 1.

N	3	5	7	9	11	13	15	17	19	21	23	25	27	29
p	2	2	3	3	2	7	3	7	3	2	5	5	3	2
k'	1	1	1	0	1	4	1	4	1	1	4	0	0	1

Table 1: q_k is irreducible whenever $k \equiv k' \pmod{p}$.

Now, following [4, Page 3154], let $\phi_k \in \mathrm{Out}(\mathbb{F}_N)$ be the outer automorphism given by:

$$a_N \mapsto a_{N-1} \mapsto a_{N-2} \mapsto \cdots \mapsto a_2 \mapsto a_1 \mapsto a_1^k a_N.$$

We use this family to conclude the remaining direction of Theorem 3:

Theorem 6. *Suppose that $k \geq 3$ is such that p_k and q_k are both irreducible. The outer automorphisms ϕ_k and ϕ_k^{-1} are non-geometric and fully irreducible. Furthermore, $\omega(\phi_k) \geq k$ while $\omega(\phi_k^{-1}) < 1 + 1/\sqrt{k}$.*

Proof. We interpret ϕ_k as a piecewise-linear homotopy equivalence f of the N–petalled rose. Since this is a positive automorphism, this is a train track map [2]. The transition matrix of this map is:

$$A := \begin{pmatrix} k & 0 & 0 & \dots & 0 & 1 \\ 1 & 0 & 0 & \dots & 0 & 0 \\ 0 & 1 & 0 & \dots & 0 & 0 \\ 0 & 0 & 1 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & 1 & 0 \end{pmatrix}$$

and, up to a sign, the characteristic polynomial of A is

$$x^N - kx^{N-1} - 1 = p_k(x).$$

We chose k so that this polynomial is irreducible and so $\omega(\phi_k) = \omega(p_k)$. Now by Lemma 4 this polynomial has $N-1$ roots that lie in the interior of the unit circle and one root in $[k, k+1]$. Hence $\lambda(\phi_k) \approx k$ and $\omega(\phi_k) = \omega(p_k) \geq k$.

On the other hand, ϕ_k^{-1} is given by:

$$a_1 \mapsto a_2 \mapsto a_3 \mapsto \cdots \mapsto a_{N-1} \mapsto a_N \mapsto a_{N-1}^{-k} a_1.$$

Again, we consider this as a piecewise-linear homotopy equivalence g of the N–petalled rose. Direct calculation of the turns involved shows that this is again a train track map and that its transition matrix is:

$$B := \begin{pmatrix} 0 & 1 & 0 & \dots & 0 & 0 \\ 0 & 0 & 1 & \dots & 0 & 0 \\ 0 & 0 & 0 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & 0 & 1 \\ 1 & 0 & 0 & \dots & k & 0 \end{pmatrix}$$

Up to a sign, the characteristic polynomial of B is

$$x^N - kx^{N-2} - 1 = q_k(x).$$

Again, we chose k so that this polynomial is irreducible and so $\omega(\phi_k) = \omega(q_k)$. Thus by Lemma 4 this polynomial has $N-2$ roots that lie in the interior of the unit circle, one root in $[-\sqrt{k+1}, -\sqrt{k-1}]$ and one root in $[\sqrt{k-1}, \sqrt{k+1}]$. Thus, $\lambda(\phi_k) \approx \sqrt{k}$ and so

$$\omega(\phi_k^{-1}) = \omega(q_k) \leq \frac{\sqrt{k+1}}{\sqrt{k-1}} \leq 1 + \frac{1}{\sqrt{k}}$$

as required.

It also follows from this computation that $\lambda(\phi_k) \neq \lambda(\phi_k^{-1})$ and so ϕ_k is a non-geometric automorphism.

Now suppose that f has no periodic Nielsen paths [10, Page 25]. Note that the local Whitehead graph of f is connected [10, Page 25] and A is a Perron–Frobenious matrix. Thus by the Full Irreducibility Criterion [10, Lemma 9.9] we have that ϕ_k is fully irreducible.

On the other hand, if f has a periodic Nielsen path then ϕ_k must be parageometric [4, Page 3155]. Thus ϕ_k^{-1} is not parageometric [5, Corollary 2] and so g has no periodic Nielsen paths. Again, the local Whitehead graph of g is connected and B is a Perron–Frobenious matrix, hence ϕ_k^{-1} is fully irreducible.

In either case, ϕ_k and ϕ_k^{-1} are both fully irreducible. □

Handel and Mosher described how there is no uniform upper bound, depending only on N, on the distance between the axes of ϕ and ϕ^{-1} [6, Section 7.1]. We finish by noting that ϕ_k is an explicit family of such outer automorphisms.

Theorem 7. *Suppose that $k \geq 5$ is such that p_k and q_k are both irreducible. There are axes for ϕ_k and ϕ_k^{-1} which are separated by at least $\left(N - \frac{5}{2}\right)\log(k) - 6$.*

Proof. For ease of notation let $\lambda_k := \lambda(\phi_k)$ and $\lambda\!\!\!^{\backprime}_k := \lambda(\phi_k^{-1})$.

Begin by noting that the Perron–Frobenious eigenvectors of the matrices A and B in the proof of Theorem 6 are

$$\begin{pmatrix} \lambda_k^{N-1} \\ \lambda_k^{N-2} \\ \vdots \\ \lambda_k \\ 1 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 1 \\ \lambda\!\!\!^{\backprime}_k \\ \vdots \\ \lambda\!\!\!^{\backprime N-2}_k \\ \lambda\!\!\!^{\backprime N-1}_k \end{pmatrix}$$

respectively. Let X_0 and Y_0 be the metric graphs in Outer space corresponding to N–petalled roses with these lengths assigned. Then X_0 and its images $X_i := X_0 \cdot \phi_k^i$ lie on an axis of ϕ_k. Similarly, Y_0 and its images $Y_i := Y_0 \cdot \phi_k^{-i}$ lie on an axis of ϕ_k^{-1}.

We note that

$$d(X_i, X_{i+1}) = \log(\lambda_k) \quad \text{and} \quad d(Y_i, Y_{i+1}) = \log(\lambda\!\!\!^{\backprime}_k).$$

Furthermore, direct calculation of the lengths of candidate loops shows that $d(X_i, Y_i) \leq d(X_i, Y_j)$ for every i and j and that

$$d(X_i, Y_i) = \log\left(\frac{\lambda_k^N - 1}{\lambda_k - 1} \cdot \frac{\lambda\!\!\!^{\backprime N}_k - \lambda\!\!\!^{\backprime N-1}_k}{\lambda\!\!\!^{\backprime N}_k - 1}\right) \geq (N-1)\log(k) - 4.$$

Therefore suppose that the distance between these two axes is realised by points X and Y. Without loss of generality we may assume that X lies in the segment $[X_i, X_{i+1}]$ and that Y lies in the segment $[Y_{j-1}, Y_j]$. However, as shown in Figure 2,

$$d(X, Y) \geq d(X_i, Y_j) - d(X_i, X) - d(Y, Y_j).$$

As $\lambda_k \leq k+1$ and $\lambda\!\!\!^{\backprime}_k \leq \sqrt{k+1}$, the right hand side of this inequality is bounded below by

$$(N-1)\log(k) - 4 - \frac{3}{2}\log(k+1) \geq \left(N - \frac{5}{2}\right)\log(k) - 6$$

as required. □



Figure 2: Axes of ϕ_k and ϕ_k^{-1}.

We note that for each ϕ_k the turns $\{a_i, a_j\}$ are all illegal. Hence these are never lone axis automorphisms [9, Theorem 3.9]. Similarly, for each ϕ_k^{-1} the turns $\{a_i^{-1}, a_{i+2}\}$ (where the indices wrap over so $a_{N+1} := a_1$ and $a_{N+2} := a_2$) are all illegal and so these are also never lone axis automorphisms.

Question 8. Are there axes of ϕ_k and ϕ_k^{-1} that remain within a uniformly bounded distance of each other?

Remark 9. From the eigenvectors above, we also see that as k goes to infinity these axes enter thinner and thinner parts of Outer space.

Acknowledgements. The author wishes to thank Yael Algom-Kfir and Christopher J. Leininger for many helpful discussions regarding this result.

References

[1] Mark C. Bell and S. Schleimer. Slow north-south dynamics on $\mathcal{PML}$. *ArXiv e-prints*, December 2015. [1, 2]

[2] Mladen Bestvina and Michael Handel. Train tracks and automorphisms of free groups. *Ann. of Math. (2)*, 135(1):1–51, 1992. [4]

[3] Benson Farb and Dan Margalit. *A primer on mapping class groups*, volume 49 of *Princeton Mathematical Series*. Princeton University Press, Princeton, NJ, 2012. [2]

[4] Michael Handel and Lee Mosher. Parageometric outer automorphisms of free groups. *Trans. Amer. Math. Soc.*, 359(7):3153–3183 (electronic), 2007. [3, 5]

[5] Michael Handel and Lee Mosher. Parageometric outer automorphisms of free groups. *Trans. Amer. Math. Soc.*, 359(7):3153–3183 (electronic), 2007. [5]

[6] Michael Handel and Lee Mosher. Axes in outer space. *Mem. Amer. Math. Soc.*, 213(1004):vi+104, 2011. [5]

[7] Serge Lang. *Fundamentals of Diophantine geometry.* Springer-Verlag, New York, 1983. [2]

[8] Gilbert Levitt and Martin Lustig. Irreducible automorphisms of F_n have north-south dynamics on compactified outer space. *J. Inst. Math. Jussieu*, 2(1):59–72, 2003. [1]

[9] L. Mosher and C. Pfaff. Lone Axes in Outer Space. *ArXiv e-prints*, November 2013. [6]

[10] C. Pfaff. Constructing and Classifying Fully Irreducible Outer Automorphisms of Free Groups. *ArXiv e-prints*, May 2012. [5]

[11] Karen Vogtmann. Automorphisms of free groups and outer space. In *Proceedings of the Conference on Geometric and Combinatorial Group Theory, Part I (Haifa, 2000)*, volume 94, pages 1–31, 2002. [1]